




# Universal Patient Care





# Universal Patient Care

## CAPACITY/REFUSAL CHECKLIST:


- Is the patient age >18 and/or emancipated minor?
- Can the patient retain and comprehend relevant information?
- Can the patient use information to make a choice?
- Is the patient NOT DANGEROUS to self or others (i.e. no suicidal or homicidal ideation)
  
- \*\*\*If all = "YES," then the patient has capacity to decline further care/transport.
- \*\*\*If any = "NO," then the patient does not have capacity to make his or her own medical decisions.
  
- Document these concepts clearly in your narrative- simply stating "alert and oriented" is not sufficient.
- Refer to Medical Policy 2 – Definition of a Patient

## Pearls

- **Recommended Exam: Minimal exam if not noted on the specific protocol is vital signs, mental status with GCS, and location of injury or complaint.**
- Any patient contact, which does not result in an EMS transport, must have a completed Patient Care Report.
- Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.
- Two complete vital sign acquisitions should occur at a minimum with any patient encounter.
- **Patient Refusal (Declining Treatment and/ or Transport):**  
Patient refusal is a high risk situation. Encourage patient to accept transport to medical facility.  
Encourage patient to allow an assessment, including vital signs. Documentation of the event is very important including a mental status assessment describing the patient's capacity to refuse care.  
**Guide to Assessing capacity:**  
**C: Patient should be able to communicate a clear choice:** This should remain stable over time. Inability to communicate a choice or an inability to express the choice consistently demonstrates incapacity.  
**R: Relevant information is understood:** Patient should be able to voice a factual understanding of the illness/ injury, the options, and the risks and benefits of recommended treatment or transport.  
**A: Appreciation of the situation:** Ability to communicate an understanding of the facts of the situation. The patient should be able to recognize the significance of the outcome potentially from their decision.  
**M: Manipulation of information in a rational manner:** Demonstrate a rational process to come to a decision. Should be able to describe the logic they are using to come to the decision, though you may not agree with decision.
- **Pediatric Patient General Considerations:**  
**A pediatric patient is defined by fitting with a Pediatric Medication/ Skill Resuscitation System, Age ≤ 15, weight ≤ 49 kg.**  
Special needs children may require continued use of Pediatric based protocols regardless of age and weight. Initial assessment should utilize the **Pediatric Assessment Triangle** which encompasses Appearance, Work of Breathing and Circulation to skin.  
The order of assessment may require alteration dependent on the developmental state of the pediatric patient. Generally the child or infant should not be separated from the caregiver unless absolutely necessary during assessment and treatment.
- Timing of transport should be based on patient's clinical condition and the agency transport policy.
- Consider consultation with Medical Control for patient(s) refusing treatment/ transport.
- Blood Pressure is defined as a Systolic/ Diastolic reading. A palpated Systolic reading may be necessary at times.
- SAMPLE: Signs/ Symptoms; Allergies; Medications; PMH; Last oral intake; Events leading to illness/ injury



# Triage





# Triage

## Pearls

- **When approaching a multiple casualty incident where resources are limited:**  
Triage decisions must be made rapidly with less time to gather information  
Emphasis shifts from ensuring the best possible outcome for an individual patient to ensuring the best possible outcome for the greatest number of patients.
- **Scene Size Up:**
  1. **Conduct a scene size up. Assure well being of responders. Determine or ensure scene safety before entering. If there are several patients with the same complaints consider HazMat, WMC or CO poisoning.**
  2. **Take Triage system kit.**
  3. **Determine number of patients. Communicate the number of patients and nature of the incident and establish incident command.**
  4. **Direct incoming resources. Identify ingress and egress path. Establish a staging area. Assign a medical officer, triage officer, transportation officer, and staging officer as personnel become available.**
- **Triage is a continual process and is a continuous process in each section as resources allow.**
- **Step 1: Global sorting:**  
Call out to those involved in the incident to walk to a designated area and assess group last.  
For those who cannot walk, have them wave/ indicate a purposeful movement and assess them second.  
Those involved who are not moving, or have an obvious life threat, assess first.
- **Step 2: Individual assessments:**  
Control major hemorrhage.  
Open airway and if child, give 2 rescue breaths.  
Perform Needle Chest Decompression Procedure if indicated.  
Administer injector antidotes if indicated.
- **Assess the first patient you encounter using the three objective criteria which can be remembered by RPM.**  
**R: Respiratory** (*Respiratory rates are difficult to measure quickly, use work of breathing and respiratory distress*)  
**P: Perfusion** (*Capillary refill can be altered by many factors including skin temperature – use age appropriate heart rates*)  
**M: Mental Status** (*Motor component of GCS score is important indicator – ability to follow commands*)
- If your patient falls into the RED TAG category, stop, place RED TAG and move on to next patient. Attempt only to correct airway problems, treat uncontrolled bleeding, or administer an antidote before moving to next patient.
- **Treatment:**  
Once casualties are triaged, a focus on treatment can begin. You may need to move patients to treatment areas.  
RED TAGs are moved/ treated first, followed by YELLOW TAGs. BLACK TAGs should remain in place.  
You may also indicate deceased patients by pulling their shirt/ clothing over their head.  
As more help arrives, then the triage/ treatment process may proceed simultaneously.
- **Lightning strike (Reverse Triage):**  
Lightning strike victims are amenable to airway, breathing, cardiac compressions as well as early defibrillation.  
Use concept of reverse triage with multiple casualties. Resuscitate lightning strikes as the priority.  
Lightning strike victims found alive do not often deteriorate quickly.
- **SMART triage tag system is utilized in NC.**



# Abdominal Pain Vomiting and Diarrhea

## History

- Age
- Time of last meal
- Last bowel movement/emesis
- Improvement or worsening with food or activity
- Duration of problem
- Other sick contacts
- Past medical history
- Past surgical history
- Medications
- Menstrual history (pregnancy)
- Travel history
- Bloody emesis / diarrhea

## Signs and Symptoms

- Pain
- Character of pain (constant, intermittent, sharp, dull, etc.)
- Distention
- Constipation
- Diarrhea
- Anorexia
- Radiation

### Associated symptoms:

Fever, headache, blurred vision, weakness, malaise, myalgias, cough, headache, dysuria, mental status changes, rash

## Differential

- CNS (increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage, vestibular)
- Myocardial infarction
- Drugs (NSAID's, antibiotics, narcotics, chemotherapy)
- GI or Renal disorders
- Diabetic ketoacidosis
- OB-Gyn disease (ovarian cyst, PID, Pregnancy)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or toxin induced
- Medication or Substance abuse
- Psychological

|          |  |
|----------|--|
|          | Consider Blood Glucose Analysis Procedure                        |
| <b>B</b> | 12 Lead ECG Procedure  |
|          | IV or IO Access Protocol UP 6                                    |
| <b>P</b> | Cardiac Monitor  |
|          | Age Appropriate Diabetic Protocol AM 2/ PM 2 <b>if indicated</b> |
|          | Pain Control Protocol UP 11 <b>if indicated</b>                  |
|          | Age Appropriate Cardiac Protocol(s) <b>if indicated</b>          |

**Age Specific Blood Pressure indicating possible shock**

**Age 0 – 28 days: SBP < 60**  
**Ages ≥ 1 month: SBP < 70**  
**Age 1 – 9: SBP < 70 + (2x Age)**

**Ages 10 – 64: SBP < 90**  
**Ages ≥ 65: SBP < 110**

**All ages Shock Index: HR > SBP**

Serious Signs/ Symptoms  
Hypotension, poor perfusion, shock

YES

NO

**A**

**Normal Saline IV TKO Or Saline Lock**

**Ondansetron 4 mg IV / IO / ODT / PO / IM**  
**Peds: 0.2 mg/kg**  
**Peds Maximum 4 mg**  
 May repeat in 15 minutes

If no response in adults  
**Compazine 5mg IM / IV / IO**  
 May repeat x 1 as needed

IV or IO Access Protocol UP 6  
Consider 2 Large Bore sites

**A**

**Normal Saline 500 mL Bolus**  
 Repeat as needed  
 Titrate SPB ≥ 90 mmHg  
**Maximum 2 L**  
**Peds: 20 mL/kg IV / IO**  
 Repeat as needed  
 Titrate to  
 Age Appropriate  
 SBP ≥ 70 + (2 x Age)  
**Maximum 60 mL/kg**

**Ondansetron 4 mg IV / IO / ODT / PO / IM**  
**Peds: 0.2 mg/kg**  
**Peds Maximum 4 mg**  
 May repeat in 15 minutes

Age Appropriate Hypotension/ Shock Protocol AM 5/ PM 3 **if indicated**

Monitor and Reassess

**Notify Destination or Contact Medical Control**

\*\* See Page Two for approved alternate Medications \*\*



# Abdominal Pain Vomiting and Diarrhea

## Prochlorperazine (Compazine)

Use with caution as it has a strong sedative effect.  
Consider combined effects with narcotic co-administration.  
Dilute in 10ml of saline- flush and administer slowly.  
May be administered first line when PT is allergic to Zofran

### Dystonic reactions:

- can occur with co-administration of psych meds.
- involuntary contractions of muscles in the extremities, face, neck, abdomen, pelvis, or larynx in either sustained or intermittent patterns that lead to abnormal movements or postures
- Treat with Benadryl 25-50mg IV/IM/IO

LR (Lactated Ringers), Normosol, Plasmalyte are acceptable substitutes for Normal Saline

**Alternative Agents:** To be used in times of drug shortage / Special Permission from Med Director

- Promethazine (Phenergan) 25mg IM x 1 - May only be given once IM (Not IV) Weight > 40kg
- Metoclopramide (Reglan) 5mg IV/IO/IM – May repeat x 1
- Droperidol 1.25mg IV - May repeat x 1 - Very sedating / may cause significant QT prolongation

## Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Abdominal/ back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and/ or lower extremity pain or diminished pulses, especially in patients over 50 and/ or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics and/ or women, especially with upper abdominal complaints.**
- **Heart Rate: Tachycardia is one of the first clinical signs of dehydration and volume depletion and typically increases as dehydration becomes more severe.**
- **Nausea without vomiting should be treated like vomiting. Patient will benefit from symptom control with antiemetic even if not actively vomiting.**
- Isolated vomiting in children is common but can be a sign of more serious pathology. Pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures) all often present with vomiting.
- Vomiting and diarrhea are common symptoms, but can be the symptoms of uncommon and serious pathology such as stroke, CO poisoning, acute MI, new onset diabetes, diabetic ketoacidosis (DKA), and organophosphate poisoning. Maintain a high index of suspicion for serious pathology.



# Altered Mental Status

## History

- Known diabetic, medic alert tag
- Drugs, drug paraphernalia
- Report of illicit drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits


## Signs and Symptoms

- Decreased mental status or lethargy
- Change in baseline mental status
- Bizarre behavior
- Hypoglycemia (cool, diaphoretic skin)
- Hyperglycemia (warm, dry skin; fruity breath; Kussmaul respirations; signs of dehydration)
- Irritability

## Differential

- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection (CNS and other)
- Thyroid (hyper / hypo)
- Shock (septic, metabolic, traumatic)
- Diabetes (hyper / hypoglycemia)
- Toxicological or Ingestion
- Acidosis / Alkalosis
- Environmental exposure
- Pulmonary (Hypoxia)
- Electrolyte abnormality
- Psychiatric disorder

|   |                                  |
|---|----------------------------------|
| Age Appropriate<br>Airway Protocol(s) AR 1, 2, 3, 5, 6<br><i>if indicated</i> |                                  |
|   | Blood Glucose Analysis Procedure |
| <b>B</b>  | 12 Lead ECG Procedure            |
|   | IV or IO Access Protocol UP 6    |
| Age Appropriate Diabetic<br>Protocol(s) AM 2/ PM 2<br><i>if indicated</i>     |                                  |





# Altered Mental Status

## Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.**
- **AMS may present as a sign of an environmental toxin or Haz-Mat exposure, protect personal safety.**
- **General:**
  - **The patient with AMS poses one of the most significant challenges.**
  - **A careful assessment of the patient, the scene, and the circumstances should be undertaken.**
  - **Assume the patient has a life threatening cause of their AMS until proven otherwise.**
  - **Pay careful attention to the head exam for signs of bruising or other injury.**
  - **Information found at the scene must be communicated to the receiving facility.**
  - **Patients not able to communicate with you coherently require a complete secondary survey (head-to-toe) exam to assess for trauma, infection, or signs of maltreatment/ abuse, or neglect.**
  - **Acute Stroke should be considered in all patients with acute AMS when < 24 hours from onset.**
- **Substance misuse:**
  - Patients ingesting substances can pose a great challenge.
  - DO NOT assume recreational drug use and/ or alcohol are the sole reasons for AMS.
  - Misuse of alcohol/ recreational drugs may lead to hypoglycemia or occult trauma.
  - More serious underlying medical and trauma conditions may be the cause.
- **Behavioral health:**
  - The behavioral health patient may present a great challenge in forming a differential.
  - DO NOT assume AMS is the result solely of an underlying psychiatric etiology.
  - Often an underlying medical or trauma condition precipitates a deterioration of a patients underlying disease.
- **Spinal Motion Restriction/ Trauma:**
  - Only utilize spinal immobilization if the situation warrants.
  - The patient with AMS may worsen with increased agitation when immobilized.
- **It is safer to assume hypoglycemia than hyperglycemia if doubt exists. Recheck blood glucose after Dextrose or Glucagon**
- Consider Restraints if necessary for patient's and/ or personnel's protection per USP 5 Restraints: Physical procedure.



# Back Pain

## History

- Age
- Past medical history
- Past surgical history
- Medications
- Onset of pain / injury
- Previous back injury
- Traumatic mechanism
- Location of pain
- Fever
- Improvement or worsening with activity


## Signs and Symptoms

- Pain (paraspinous, spinous process)
- Swelling
- Pain with range of motion
- Extremity weakness
- Extremity numbness
- Shooting pain into an extremity
- Bowel / bladder dysfunction

## Differential

- Muscle spasm / strain
- Herniated disc with nerve compression
- Sciatica
- Spine fracture
- Kidney stone
- Pyelonephritis
- Aneurysm
- Pneumonia
- Spinal Epidural Abscess
- Metastatic Cancer
- AAA

|          |   |
|----------|---|
| <b>B</b> | Consider Cardiac Etiology<br>12 Lead ECG Procedure<br><i>if indicated</i> |
| <b>P</b> | Cardiac Monitor<br><i>if indicated</i>                                    |
|          | Age Appropriate<br>Cardiac Protocol(s)<br><i>if indicated</i>             |





# Back Pain

Consider Abdominal Aortic Aneurysm with severe back pain. Classic presentation is abdominal pain radiating to the back. A pulsatile mass may be felt on thin patients.

Avoid Toradol in these patients

Consider Thoracic Aortic Dissection with severe upper back pain - Pain between the shoulder blades and chest pain.

## Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Neuro, Lower extremity perfusion, Back**
- **Back pain is one of the most common complaints in medicine and affects more than 90% of adults at some point in their life. Back pain is also common in the pediatric population. Most often it is a benign process but in some circumstances can be life or limb threatening.**
- **Consider pregnancy or ectopic pregnancy with abdominal or back pain in women of childbearing age.**
- **Consider abdominal aortic aneurysm with abdominal pain especially in patients over 50 and/ or patients with shock/ poor perfusion. Patients may have abdominal pain and/ or lower extremity pain with diminished pulses. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics and/ or women especially with upper abdominal complaints.**
- **Red Flags which may signal a more serious process associated with back pain:**
  - Age > 50 or < 18
  - Neurological deficit (leg weakness, urinary retention, or bowel incontinence)
  - IV Drug use
  - Fever
  - History of cancer, either current or remote
  - Night time pain in pediatric patients
- **Cauda equina syndrome is where the terminal nerves of spinal cord are being compressed (Symptoms include):**
  - Saddle anesthesia (numbness between the genitalia and rectum)
  - Recent onset of bladder and bowel dysfunction. (Urine retention and bowel incontinence)
  - Severe or progressive neurological deficit in the lower extremity.
  - Motor weakness of thigh muscles or foot drop
- **Back pain associated with infection:**
  - Fever/ chills.
  - IV Drug user (consider spinal infection)
  - Recent bacterial infection like pneumonia.
  - Immune suppression such as HIV or patients on chronic steroids like prednisone.
  - Meningitis.
- **Spinal motion restriction in patients with underlying spinal deformity should be maintained in their functional position.**
- **Kidney stones typically present with an acute onset of flank pain which radiates around to the groin area.**



# IV or IO Access

## History


- Chronic medical conditions requiring recurrent need for IV access for medication, hydration, or blood sampling.
- Medical condition requiring administration of IV medications at home.
- End-stage renal disease requiring hemodialysis.
- Chronic medical condition requiring IV nutrition.

## Signs and Symptoms

- Fever
- Bleeding
- Hypotension
- Redness, swelling, and/or pain at IV catheter site
- Shortness of breath
- Chest pain
- IV catheter patency

## Differential

- Infection or sepsis
- Infection of catheter
- Clotted IV catheter
- Air embolism
- Pneumothorax
- Overdose of home medication
- Shock



## Pearls

- **Frequent encounter of patients with IV access devices and confusion as to which device can be accessed and used by EMS providers are common.**
- **If unclear about device use, always ask "Is this device used for dialysis?"**
- **When accessing central catheter, always ensure sterility of catheter connection point by cleaning port with alcohol, or similar disinfectant, 2 – 3 times prior to access.**
- **Central line catheters placed for administration of chemotherapy, medications, electrolytes, antibiotics, and blood are available to EMS providers for access and administration of fluids, medications, antibiotics, and blood products.**
- **Central line catheters placed for hemodialysis are NOT available for access by EMS providers unless the patient is in cardiac arrest.**
- Long term IV access is frequently needed for a variety of indications:
  - Medication administration such as antibiotics, pain relief, or chemotherapy.
  - Administration of IV nutrition or feeding.
  - Need for multiple IV line access or recurrent blood sampling.
  - Poor vasculature requiring repeated attempts at IV access.
  - End-stage renal disease requiring hemodialysis.
- Common complications of central access devices:
  - Infection
  - Damage to vasculature
  - Air embolism
  - Loss of patency due to clogging or clotting
  - Pneumothorax

# IV or IO Access

## Types of IV catheters:


### **Port-a-Cath® :**

Surgically implanted device allowing easy access to venous system. The port and the catheter are all placed beneath the skin. Requires a special kit and a specific needle to access.

**Paramedic does NOT routinely access this device.**

**Paramedic may utilize if already accessed with needle/ extension.**

**Paramedic may access if trained on procedure with proper equipment available.**



### **Dialysis Catheter:**

Surgically implanted device used to access the vasculature for hemodialysis.

May be tunneled under the skin with access on outside of skin surface or may be non-tunneled with greater portion of catheter on outside of skin surface.

Catheter has a RED port indicating use for dialysis:


Most catheters have a RED port and a BLUE port.

Some catheters have a RED port and a WHITE port.

Dialysis catheters may be used for both short and long-term dialysis and should not be accessed or used for delivery of fluids, medications, antibiotics, or blood products as it increases risk of infection, which then requires removal and subsequent loss of dialysis access.

**Paramedic and AEMT do NOT routinely access this device.**

**Paramedic and AEMT MAY access during cardiac arrest only (Only if IV or IO access cannot be established.)**



### **PICC (Peripherally Inserted Central Catheters):**


Long catheter inserted into a vein in arm or leg (less common) with the tip of the catheter positioned into the central circulation.

Used for long-term IV fluids, medication administration, blood administration or blood draws.

May have 1 or 2 ports (possibly more, but less common.)

Port ends usually white, blue, or purple. (May be red, less common and is not used for dialysis.)

**Paramedic and AEMT may access and utilize following clean technique.**



### **Central Lines:**


Catheter placed in large vein in the neck, under the clavicle, or in the groin.

Used for long-term IV fluids, medication administration, blood administration or blood draws.

May have 1 - 4 ports (possibly more, but less common.)

Port ends usually white, blue, or purple. (May be red, less common and is not used for dialysis.)

**Paramedic and AEMT may access and utilize following sterile technique.**





# Dental Problems

## History


- Age
- Past medical history
- Medications
- Onset of pain / injury
- Trauma with "knocked out" tooth
- Location of tooth
- Whole vs. partial tooth injury

## Signs and Symptoms

- Bleeding
- Pain
- Fever
- Swelling
- Tooth missing or fractured

## Differential

- Decay
- Infection
- Fracture
- Avulsion
- Abscess
- Facial cellulitis
- Impacted tooth (wisdom)
- TMJ syndrome
- Myocardial infarction





# Dental Problems

**\*\* May utilize TXA soaked gauze or nebulized for severe oral pharyngeal bleeding \*\***

## Pearls

### **Procedure:**

- **Apply 10 ml of a 1 g/10 ml Tranexamic acid (TXA) solution to a gauze pad or cotton**
- **Pack tranexamic acid-soaked gauze or cotton into the bleeding tooth socket and apply direct pressure with patient closing teeth to exert pressure**
- **Gauze may be removed after 20 minutes, and bleeding has stopped.**
  
- **Recommended Exam: Mental Status, HEENT, Neck, Chest, Lungs, Neuro**
- Significant soft tissue swelling to the face or oral cavity can represent a cellulitis or abscess.
- Scene and transport times should be minimized in complete tooth avulsions. Reimplantation is possible within 4 hours if the tooth is properly cared for, but unlikely when > 1 hour from time of injury.
- Cardiac chest pain may radiate to the jaw and teeth mimicking dental pain.
- **Avulsed tooth:**
  - Handle tooth by the crown, do not touch the root.
  - Rinse tooth if soiled but do not scrub, as this can damage the ligaments vital for possible reimplantation.
  - Rinse with mild, commercial tooth solution, normal saline or lactated ringers, or the patient's own saliva if dry.
  - Transport tooth in milk, commercial solution, patient's own saliva, or IV solution in a container to protect.
- **TXA Use in Dental Bleeding:**
  - May be used topically or nebulized for severe oropharyngeal bleeding
  - TXA offers modest benefit as TXA instilled gauze combined with direct pressure.



# Emergencies Involving Indwelling Central Lines

## History


- Central Venous Catheter Type  
Tunneled Catheter (Broviac/ Hickman)
- PICC (peripherally inserted central catheter)
- Implanted catheter (Mediport/ Hickman)
- Occlusion of line
- Complete or partial dislodge
- Complete or partial disruption

## Signs and Symptoms

- External catheter dislodgement
- Complete catheter dislodgement
- Damaged catheter
- Bleeding at catheter site
- Internal bleeding
- Blood clot
- Air embolus
- Erythema, warmth or drainage about catheter site indicating infection

## Differential

- Fever
- Hemorrhage
- Reactions from home nutrient or medication
- Respiratory distress
- Shock



**Use Sterile Technique:**

- During manipulation of central line
- When accessing central line

## Pearls

- **Always involve family/ caregivers as they may have specific knowledge and skills related to catheter device.**
- **Use strict sterile technique when accessing/ manipulating an indwelling catheter.**
- **Cardiac arrest: May access central catheter and utilize if functioning properly.**
- Do not attempt to force catheter open if occlusion evident.
- Some infusions may be detrimental to stop. Ask family or caregiver if it is appropriate to stop or change infusion.
- Hyperalimentation infusions (IV nutrition): If stopped for any reason, monitor for hypoglycemia.



# Epistaxis

## History


- Age
- Past medical history
- Medications (HTN, anticoagulants, aspirin, NSAIDs)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

## Signs and Symptoms

- Bleeding from nasal passage
- Pain
- Nausea
- Vomiting

## Differential

- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension



### Age Specific Blood Pressure indicating possible shock

Age 0 – 28 days: SBP < 60  
 Ages ≥ 1 month: SBP < 70  
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90  
 Ages ≥ 65: SBP < 110

All ages Shock Index:  
 HR > SBP

**A** May soak RHINO ROCKET in TXA prior to insertion

**\*\* Reserve hemostatic devices for severe bleeds not stopped by traditional means \*\***



# Epistaxis

## Pearls

### **Procedure:**

- Apply 10 ml of a 1 g/10 ml Tranexamic acid (TXA) solution to a gauze pad
- Insert tranexamic acid-soaked gauze into nasal cavity
- Gauze may be removed after 20 minutes, and bleeding has stopped.
- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Neuro**
- **Tranexamic acid (TXA) Use in Epistaxis:**
  - May be used topically or nebulizer
  - No clear evidence that Tranexamic acid (TXA) provides benefit over conventional vasoconstrictors and sustained direct pressure.
- It is very difficult to quantify the amount of blood loss with epistaxis.
- Bleeding may also be occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Anticoagulants include warfarin (Coumadin), Apixaban (Eliquis), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), and many over the counter headache relief powders.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/ dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.



# Fever/ Infection Control

## History

- Age
- Duration of fever
- Severity of fever
- Past medical history
- Medications
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Environmental exposure
- Last acetaminophen or ibuprofen

## Signs and Symptoms

- Warm
- Flushed
- Sweaty
- Chills/Rigors


### Associated Symptoms (Helpful to localize source)

- Myalgias, cough, chest pain, headache, dysuria, abdominal pain, mental status changes, rash

## Differential

- Infections / Sepsis
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease
  - Arthritis
  - Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis

|          |  |
|----------|--|
|          | Contact, Droplet, and Airborne Precautions<br>See Pearls |
| <b>B</b> | Temperature Measurement Procedure<br><i>if available</i> |
|          | IV or IO Access Protocol UP 6<br><i>if indicated</i>     |



**\*\* Do not give if PO or PR dose already given within the past 4 hours\*\***  
(May be given via IO only if IO is ALREADY in place for patient care)

## Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Febrile seizures are more likely in children with a history of febrile seizures and with a rapid elevation in temperature.
- Patients with a history of liver failure should not receive acetaminophen.
- Rehydration with fluids increases the patient's ability to sweat and improves heat loss.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen not acetaminophen.
- NSAIDs should not be used in the setting of environmental heat emergencies.
- **Do not** give aspirin to a child, age ≤ 15 years.
- Manufacturer recommends to be infused over 15 minutes. This recommendation is related to the formulation propacetamol, the prodrug to acetaminophen, approved in Europe, which caused pain on infusion, and data from the clinical development of acetaminophen. The formulation of IV acetaminophen is not a prodrug and does not have the same adverse reaction. An objective chart review study has shown the lack of side effects of rapidly infusing intravenous acetaminophen. There it is acceptable to infuse IV Acetaminophen over 3 to 4 minutes in adults only. Pediatrics has not been studied.
- Agency Medical Director does not require contact of medical control prior to EMT/ EMR administering any medication.



# Pain Control

## History

- Age
- Location
- Duration
- Severity (1 - 10)
- If child use Wong-Baker faces scale
- Past medical history
- Medications
- Drug allergies

## Signs and Symptoms

- Severity (pain scale)
- Quality (sharp, dull, etc.)
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

## Differential

- Per the specific protocol
- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural/ Respiratory
- Neurogenic
- Renal (colic)



## Specific Complaint Protocol

Assess Pain Severity

Combination of Pain Scale, MOI, circumstances, Injury or Illness severity

Mild < 6/10

Moderate to Severe > 6/10

|  |   |
|--|---|
| <b>B</b>   | <b>Ibuprofen 10 mg/kg PO</b><br>(400 – 600 mg typical adult)<br>Maximum 800 mg  |
|  | Or<br><b>Acetaminophen 15 mg/kg PO</b><br>(325 – 1000 mg typical adult)<br>Maximum 1000 mg                            |
| <b>Nitrous Oxide 70:30 to 50:50 Mix NO/O<sub>2</sub></b>                                       |   |
| IV or IO Access Protocol UP 6  |   |
| <b>A</b>   | <b>Ketorolac 15 mg IV / IO</b><br>30 mg IM<br>Peds: 0.5 mg/kg IV / IO / IM<br>Maximum 30 mg                           |
|  | <b>IV APAP – If not already given PO</b><br>Acetaminophen 1g IV/IO x 1<br>Peds: 15mg/kg IV over 10 min<br>Max: 1 gram |
| Monitor and Reassess<br>Every 10 minutes following medication                                  |   |
| If no improvement reassess pain<br>Consider Paramedic Level Medications<br><i>if indicated</i> |   |

|  |   |
|--|---|
| <b>P</b>   | IV or IO Access Protocol UP 6<br>Cardiac Monitor  |
|  | <b>Fentanyl 50 – 100 mcg IV / IO / IM / IN</b><br>Repeat 25 mcg every 10 minutes as needed<br>Maximum 300 mcg   |
|  | <b>Adult &amp; Ped (&gt;6 mo): 1 mcg / kg IV / IO / IM / IN</b><br>Repeat 0.5 mcg / kg every 10 minutes prn<br>Max single dose: 100 mcg<br>Adult Maximum: 300 mcg<br>Pediatric Maximum: 100 mcg |
| OR   |   |
| <b>Morphine 2 – 4 mg IV / IO / IM</b><br>Repeat every 5 minutes as needed<br>Maximum 20 mg   |   |
| <b>Ketamine 0.3 mg/kg IV / IO</b><br>Infuse or IV push over 10 minutes<br>May repeat every 20 minutes<br>Maximum 30 mg single dose |   |
| Maximum 3 total doses<br>Or<br>Ketamine 1 mg/kg IN<br>Maximum 1 total dose   |   |



Notify Destination or  
Contact Medical Control





# Pain Control

Fentanyl dosing in adults is at the provider discretion on which method to use – weight based or aliquot methods

Ketamine is a 2<sup>nd</sup> line option after one dose of opioid has been given.

**Nitrous Oxide** is a nonscheduled medical gas. Nonparamedic services wishing to utilize Nitrous Oxide will need a security plan and a method to track usage in PSI, Weight(grams), or minutes administered per use. Nitrous does not require a controlled substance license

(State or Federal DEA is not necessary)

Must be approved by the medical director

**Alternative Medications for Pain:** Hydromorphone Adult: 0.5 mg IV / IO / IM -- Repeat every 10 minutes as

## Pearls

- **Recommended Exam: Mental Status, Area of Pain, Neuro**
- **Pain severity (0-10) is a vital sign to be recorded before and after PO, IV, IO or IM medication delivery and at patient hand off. Monitor BP closely as sedative and pain control agents may cause hypotension.**
- **Ketamine:**  
**Ketamine may be used in patients who are outside a Pediatric Medication/ Skill Resuscitation System product.**  
**Ketamine may be used in patients who fit within a Pediatric Medication/ Skill Resuscitation System product only with DIRECT ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.**
- **Ketamine: appropriate indications for pain control:**  
Patients who have developed opioid-tolerance. Sickle cell crisis patients with opioid-tolerance.  
Patients who have obstructive sleep apnea.  
May use in combination with opioids to limit total amount of opioid administration.
- **Ketamine: caution when using for pain control:**  
Slow infusion or IV push over 10 minutes is associated with less side effects. Do not administer by rapid IV push.  
Avoid in patients who have cardiac disease or uncontrolled hypertension.  
Avoid in patients with increased intraocular pressure such as glaucoma.  
Avoid use in combination with benzodiazepines due to depressed respiratory drive.
- **Both arms of the treatment protocol may be used in concert. For patients in Moderate pain for instance, you may use the combination of an oral medication and parenteral if no contraindications are present.**
- **Pediatrics:**  
For children use Wong-Baker faces scale or the FLACC score (see Assessment Pain Procedure ASP 2)  
Use Numeric (> 9 yrs), Wong-Baker faces (4-16yrs) or FLACC scale (0-7 yrs) as needed to assess pain.
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- All patients who receive IM or IV medications must be observed 15 minutes for drug reaction in the event no transport occurs.
- Do not administer **Acetaminophen** to patients with a history of liver disease.
- Burn patients may required higher than usual opioid doses to titrate adequate pain control.
- Consider agency-specific anti-emetic(s) for nausea and/ or vomiting.



# Police Custody

## History


- Traumatic Injury
- Drug Abuse
- Cardiac History
- History of Asthma
- Psychiatric History

## Signs and Symptoms

- External signs of trauma
- Palpitations
- Shortness of breath
- Wheezing
- Altered Mental Status
- Intoxication/Substance Abuse

## Differential

- Agitated Delirium Secondary to Psychiatric Illness
- Agitated Delirium Secondary to Substance Abuse
- Traumatic Injury
- Closed Head Injury
- Asthma Exacerbation
- Cardiac Dysrhythmia





# Police Custody

- **Pearls**

- **Patient does not have to be in police custody or under arrest to utilize this protocol.**
- **Local EMS agencies should formulate a policy with local law enforcement agencies concerning patients requiring EMS and Law Enforcement services simultaneously.**
- **Agencies should work together to formulate a disposition in the best interest of the patient.**
- **Patients restrained by law enforcement devices must be transported and accompanied by a law enforcement officer in the patient compartment who is capable of removing the devices. However, when rescuers have utilized restraints in accordance with Restraint Procedure, the law enforcement agent may follow the ambulance during transport.**
- **All patients who receive either physical and chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
- The responsibility for patient care rests with the highest authorized medical provider on scene per North Carolina law.
- If an asthmatic patient is exposed to irritant/ pepper spray and released to law enforcement, all parties should be advised to immediately contact EMS if wheezing/ difficulty breathing occurs.
- All patients with decision-making capacity in police custody retain the right to participate in decision-making regarding their care and may request care or refuse care of EMS.
- If extremity/ chemical/ law enforcement restraints are applied, follow USP 5 Restraints: Physical.
- **Consider Haldol or Droperidol for patients with history of psychosis or a benzodiazepine for patients with presumed substance misuse.**
- **Haldol is acceptable treatment in pediatric patients  $\geq 12$  years old. Safety and efficacy is not established in younger ages. Contact Medical Control for advice as needed.**
- **Hyperactive Delirium with Severe Agitation:**
  - Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia and increased strength.
  - Potentially life-threatening and associated with use of physical control measures, including physical restraints and Tasers.
  - Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents. Alcohol withdrawal or head trauma may also contribute to the condition.
- **If patient suspected of Hyperactive Delirium with Severe Agitation suffers cardiac arrest, consider a fluid bolus, administration of calcium gluconate (or chloride), and sodium bicarbonate early.**
- Do not position or transport any restrained patient in such a way that could impact the patient's respiratory or circulatory status.
- **Pepper Spray (Oleoresin Capsicum)/ CS Gas (Tear Gas) Exposure Care:**
- Be aware of cross contamination when treating patients
- Severe complications are possible with the following patients:
  - Elderly
  - Cardiac
  - COPD
  - Asthma
- Flush the affected eye(s) with normal saline - Be careful not to flush into an unaffected eye
- Capsicum exposure can also be neutralized with commercial wipes or spray
- Always wear gloves & eye protection when flushing contaminated patients
- Patients exposed to chemical spray, with or without history of respiratory disease, may develop respiratory complaints up to 20 minutes post exposure.



# Seizure

## History

- Reported / witnessed seizure activity
- Previous seizure history
- Medical alert tag information
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Document number of seizures
- Alcohol use, abuse or abrupt cessation
- Fever

## Signs and Symptoms

- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconscious

## Differential

- CNS (Head) trauma
- Tumor
- Metabolic, Hepatic, or Renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, Medications, Non-compliance
- Infection / Fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypoglycemia

|   |
|---|
| Age Appropriate<br>Airway Protocol(s) AR 1, 2, 3, 5, 6<br><b>as indicated</b>                 |
| Altered Mental Status<br>Protocol UP 4<br><b>if indicated</b>                                 |
| Childbirth/ Labor Protocol AO 1<br>Obstetrical Emergency Protocol AO 3<br><b>if indicated</b> |
| Behavioral Protocol UP 17, 18, 19<br><b>if indicated</b>                                      |
| Loosen any constrictive clothing<br>Protect patient   |

Active Seizure Activity

IV / IO Access

NO

NO

|          |   |
|----------|---|
| <b>P</b> | <b>≥ 49 kg</b><br><b>Midazolam 5 - 10 mg IM / IN</b><br>May repeat in 5 minutes if needed<br><b>Maximum 20 mg</b>                                       |
|          | <b>&lt; 49 kg</b><br><b>Midazolam 0.2 mg/kg IM / IN</b><br><b>Maximum single dose 5 mg</b><br>May repeat in 5 minutes if needed<br><b>Maximum 10 mg</b> |

|          |  |
|----------|--|
| <b>P</b> | <b>Midazolam 2 – 2.5 mg IV / IO</b><br>May repeat every 3 to 5 minutes as needed<br><b>Maximum 10 mg</b>   |
|          | <b>Peds: Midazolam 0.2 mg/kg IV / IO</b><br><b>Maximum single dose 2.5 mg</b><br>May repeat every 3 to 5 minutes as needed<br><b>Maximum 10 mg</b>   |
| <b>P</b> | For seizures refractory to Benzo's:<br>Minimum: Two Benzo Doses<br><br><b>Ketamine 1mg/kg IV/IO x 1</b><br><br>Or<br><br><b>Levetiracetam (Keppra) 1 gram IV/IO x 1</b><br><br>Peds: 15 mg/kg of mixture over 5 minutes<br>Max 1 gram<br>** If refractory after 5 minutes give both ** |

|   |  |  |
|---|--|--|
| Blood Glucose Analysis Procedure  |  |  |
| IV or IO Access Protocol UP 6<br><b>if indicated</b>  |  |  |
| <table border="1"> <tr> <td style="background-color: #4a86e8; color: white; text-align: center; vertical-align: middle;"><b>P</b></td> <td>           Cardiac Monitor<br/> <b>if indicated</b> </td> </tr> </table> | <b>P</b>                               | Cardiac Monitor<br><b>if indicated</b> |
| <b>P</b>  | Cardiac Monitor<br><b>if indicated</b> |  |
| Monitor and Reassess  |  |  |

**Notify Destination or Contact Medical Control**



# Seizure

## Alternative Seizure Medications

- Lorazepam 1-2 mg IV only (due to long onset)
- May repeat in 3-5 minutes Max Dose 4mg total
- Diazepam 5mg IV / IO Adults, Peds 0.1 - 0.3 mg/kg, Ped Rectal 0.5 mg/kg
  
- Ketamine 1mg/kg IV / IO x 1 -- use only if allergic to Benzodiazepines OR active seizures greater than 5 minutes refractory to Benzodiazepines – two doses
  
- Keppra is produced in 100mg/ml vials or several premixed forms – Keppra is safe for rapid infusion over 5 minutes maximum. Rapid infusion is safe and preferred in status epilepticus.

Active seizure in known or suspected pregnancy greater than 20 weeks, administer Magnesium Sulfate 2g IV/IO over 2-3 minutes, may repeat dose x1.

### Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **Brief seizure-like activity can be seen following ventricular fibrillation or ventricular tachycardia associated cardiac arrest.**
- **Status epilepticus is defined by seizure activity lasting > 5 minutes or multiple seizures without return to baseline.**
- **Most seizure activity is brief, lasting only 1 – 2 minutes, and is associated with transient hypoventilation.**
- **Be prepared for airway problems and continued seizures.**
- **Seizure activity may be a marker of closed head injury, especially in the very young, examine for trauma.**
- **Adult:**
  - **Midazolam 10 mg IM is effective in termination of seizures.**
  - **Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.**
- **Pediatrics:**
  - **Midazolam 0.2 mg/kg (Maximum 5 mg) IM is effective in termination of seizures.**
  - **Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.**
- **Do not delay administration of anti-epileptic drugs to check for blood glucose.**
- **Grand mal seizures (generalized)** are associated with loss of consciousness, incontinence, and tongue trauma.
- **Focal seizures** affect only a part of the body and are not usually associated with a loss of consciousness, but can propagate to generalized seizures with loss of consciousness.
- **Be prepared to assist ventilations especially if diazepam or midazolam is used.**
- **For any seizure in a pregnant patient, follow the AO 3 Obstetrical Emergencies Protocol.**
- **Midazolam (Versed) is shown to be as effective with IM route as Lorazepam (Ativan) is via the IV or IO route.**
- **Lorazepam (Ativan) is not as effective when administered IM. IV or IO route is preferred.**
- **Diazepam (Valium) is not effective when administered IM. Give IV or Rectally.**
- **Optimal conditions for patients refusing transport following a seizure:**

|   |  |
|---|--|
| Known history of seizures/epilepsy            | Seizure not associated with drugs or alcohol |
| Full recovery to baseline mental status       | Only 1 seizure episode in the past hour      |
| No injuries requiring treatment or evaluation | Seizure not associated with pregnancy        |
| Adequate supervision                          |  |



# Suspected Stroke

## History


- Previous CVA, TIA's
- Previous cardiac / vascular surgery
- Associated diseases: diabetes, hypertension, CAD
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma
- Sickle Cell Disease
- Immune disorders
- Congenital heart defects
- Maternal infection / hypertension

## Signs and Symptoms

- Altered mental status
- Weakness / Paralysis
- Blindness or other sensory loss
- Aphasia / Dysarthria
- Syncope
- Vertigo / Dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension / hypotension

## Differential

- See Altered Mental Status
- TIA (Transient ischemic attack)
- Seizure
- Todd's Paralysis
- Hypoglycemia
- Stroke
  - Thrombotic or Embolic (~85%)
  - Hemorrhagic (~15%)
- Tumor
- Trauma
- Dialysis / Renal Failure



**Transport Destination Facility based on:**  
**Stroke**  
**EMS Triage and Destination Plan**

- Refer to plan for Last Known Well Times and Stroke Severity Assessment Values
- Combination of factors will drive transport destination decision-making based on your local region

**Transport based on:**  
**STROKE**  
**EMS Triage and Destination Plan**  
**Keep Scene Time to ≤ 15 Minutes**  
**Provide Early Notification of Destination Facility**  
**Suspected Stroke Alert**

|          |  |
|----------|--|
|          | Reperfusion Checklist Procedure<br><i>if time allows completion</i>                                    |
| <b>B</b> | Cardiac Monitor  |
|          | IV or IO Access Protocol UP 6<br>Preferably 2 Sites<br><i>See Pearls</i>                               |
| <b>A</b> | Venous Access Blood Draw Procedure<br><i>if applicable</i>   |
|          | Cardiac Monitor  |
| <b>P</b> | <b>Persistent Hypertension:</b><br>If SBP ≥ 185 or DBP ≥ 110 mmHg<br>After 3 readings, 5 minutes apart |
|          | Contact Receiving Facility for guidance on hypertension treatment                                      |



# Suspected Stroke

Utilize RACE Scale EMS Stroke Assessment Tool

All patients with a positive stroke screen with symptoms less than 24 hours are time critical

If LKWT is less than 4.5 hours - pt's are candidates for thrombolytics if they qualify. Transport to nearest stroke capable hospital.

If LKWT is 4.5hrs to 24hrs with NEW SEVERE UNILATERAL DEFICITS.

Please transport or fly the patient to the nearest comprehensive stroke center to be evaluated by interventional stroke team. CEMC is acceptable if timing unclear, aeromedical unavailable, or operational limitations preventing transport to comprehensive stroke center

If transporting to CarolinaEast (CEMC) and symptoms less than 24 hours – please utilize Pulsara for a STROKE ALERT.

## Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used in the EMS Acute Stroke Care Toolkit.**
- **Acute Stroke care is evolving rapidly. Time of Onset/ Last Seen Normal may be changed at any time depending on the capabilities and resources of your regional hospital(s).**
- **Refer to your Stroke: EMS Triage and Destination Plan which should be updated when community resources change.**
- **Time of Onset or Last Seen Normal:**
  - **One of the most important items the pre-hospital provider can obtain, of which all treatment decisions are based.**
  - **Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT “about 45 minutes ago.”)**
  - **Without this information patient may not be able to receive thrombolytics at facility.**
  - **Wake up stroke: Time starts when patient was last awake or symptom free. Likely Bedtime.**
- **Time of Symptom Discovery:**
  - **Time when symptoms of stroke are first noticed by patient, bystanders, witnesses, or family/ caregivers.**
- **Sources of information pertaining to Last Known Well Time or Symptoms Onset:**
  - **You are often in the best position to determine the actual Time of Onset while you have family, friends or caretakers available.**
  - **Often these sources of information may arrive well after you have delivered the patient to the hospital. Delays in decisions due to lack of information may negatively impact patient care.**
  - **Obtain contact information (phone number and name) of best witnesses and give to hospital providers.**
- **The **Reperfusion Checklist** should be completed for any suspected stroke patient as time allows.**
- **If possible place 2 IV sites, preferably above the wrists, and if possible both in the left upper extremity.**
- **Blood Draw:**
  - **Many stroke centers utilize EMS venous blood samples. Follow your local policy and procedures.**
  - **The differential listed in the UP 4 Altered Mental Status Protocol should also be considered.**
  - **Be alert for airway problems (swallowing difficulty, vomiting/aspiration).**
  - **Hypoglycemia can present as a localized neurologic deficit, especially in the elderly.**
  - **Document the EMS Stroke Screen, Stroke Severity Score, and Stroke Alert notification time in the ePCR or PCR.**
  - **Agencies may use validated pre-hospital stroke screen of choice.**
- **Pediatrics:**
  - **Strokes do occur in children, they are slightly more common in ages < 2, in boys, and in African-Americans.**
  - **Newborn and infant symptoms consist of seizures, extreme sleepiness, and using only one side of the body.**
  - **Children and teenagers symptoms may consist of severe headaches, vomiting, sleepiness, dizziness, and/or loss of balance or coordination.**



# Suspected Sepsis

## History

- Duration and severity of fever
- Past medical history
- Medications/ Recent antibiotics
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Indwelling medical device
- Last acetaminophen or ibuprofen
- Recent Hospital/ healthcare facility
- Bedridden or immobile
- Elderly and very young – at risk
- Prosthetic device / indwelling device

## Signs and Symptoms

- Warm
- Flushed
- Sweaty
- Chills/ Rigors
- Delayed cap refill
- Mental status changes

### Associated Symptoms (Helpful to localize source)

- myalgias, cough, chest pain, headache, dysuria, abdominal pain, rash

## Differential

- Infections: UTI, Pneumonia, skin/ wound
- Cancer/ Tumors/ Lymphomas
- Medication or drug reaction
- Connective tissue disease: Arthritis, Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis
- Hypoglycemia/hypothermia
- MI/ CVA

|  |   |
|--|---|
| Consider: Contact, Droplet, and Airborne Precautions     |   |
| Temperature Measurement Procedure<br><i>if available</i> |   |
|  | Fever/ Infection Control Protocol UP 10<br><i>if needed</i> |
|  | Altered Mental Status Protocol UP 4<br><i>if needed</i>     |
| <b>B</b>   | 12 Lead ECG Procedure                                       |
|  | IV or IO Access Protocol UP 6<br><i>If indicated</i>        |
| <b>P</b>   | Cardiac Monitor   |


**Age Specific Blood Pressure indicating possible shock**

Age 0 – 28 days: SBP < 60  
 Ages ≥ 1 month: SBP < 70  
 Age 1 – 9: SBP < 70 + (2x Age)

Ages 10 – 64: SBP < 90  
 Ages ≥ 65: SBP < 110

All ages Shock Index:  
 HR > SBP

Exit to  
 Age Appropriate  
 Condition Appropriate  
 Protocol(s)



**SEPSIS ALERT**  
 Notify Receiving Facility Immediately

**MAP**  
 (Mean Arterial Pressure)

**SBP + 2(DBP)**  
 3

Monitor usually calculates this value on screen

**Adult SIRS Criteria**  
 Temperature  
 ≥ 100.4° F (38° C)  
 Or  
 ≤ 96.8° F (36° C)  
**AND**  
 Any 1 of the following:  
 HR > 90  
 RR > 20  
 EtCO < 25 mmHg

**Adult qSOFA Criteria**  
 SBP ≤ 100 mmHG  
 RR ≥ 22  
 AMS or new mental status change

**Pediatrics SIRS Criteria**  
 Temperature  
 Same as adult

**AND**  
 Heart Rate  
 1 month – 1 year > 180  
 2 – 5 years > 140  
 6 – 12 years > 130  
 13 – 18 years > 120

**A**

Venous Access Blood Draw  
*if applicable*

**Normal Saline 500 mL Bolus**  
 Repeat as needed  
 Titrate SPB ≥ 90 mmHg  
 MAP > 65 mmHg  
**Maximum 2 L**

**Peds: 20 mL/kg IV / IO**  
 Repeat to titrate  
 Age Appropriate  
 SBP ≥ 70 + 2 x Age  
**Maximum 60 mL/kg**

Draw Blood Cultures if equipped/  
 trained

**Rocephin 1g IV/IO x 1**  
 Peds: 50mg/kg IV/IO up to 1gm  
 over 10 minutes  
 See Page 2 \*\*

Age Appropriate  
 Hypotension/ Shock  
 Protocol AM 5/ PM 3

**Notify Destination or Contact Medical Control**



# Suspected Sepsis

\*\* Organizations may utilize Zosyn 3.375g or 4.5g IV OR Cefepime 2g IV as an alternative for Sepsis also \*\*

One Antibiotic is required

Rocephin may be used for patients with mild allergy to penicillin.

Avoid in severe allergies and anaphylaxis - Avoid when allergic to Cephalosporins - Keflex, Omnicef, Cefdinir, Cefuroxime

If less than age 18 - Rocephin dosage 50mg/kg - Max 1g over 10 minutes (May slow push adults)

Blood Cultures using Steripath is highly recommended but not required for nonhospital agencies.

If transporting to CarolinaEast – use Pulsara to activate a Sepsis Alert

## Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Recommended Exam Pediatrics: In childhood, physical assessment reveals important clues for sepsis. Look for mental status abnormalities such as anxiety, restlessness, agitation, irritability, confusion, or lethargy. Cardiovascular findings to look for include cool distal extremities, capillary refill >3 seconds, or mottled skin.**
- **Sepsis is a life threatening condition where the body's immune response to infection injures its own tissues and organs.**
- **Severe sepsis is a suspected infection with 2 or more SIRS criteria (or qSOFA) along with organ dysfunction, such as AMS, hypotension, or hypoxia.**
- **Septic shock is severe sepsis and poor perfusion unimproved after fluid bolus.**
- **Agencies administering antibiotics should inquire about drug allergies specific to antibiotics or family of antibiotics.**
- **Following each fluid bolus, assess for pulmonary edema. Consider administration of agency specific vasopressor.**
- **Supplemental oxygen should be given and titrated to oxygenation saturation  $\geq 92\%$ .**
- **EKG should be obtained with suspected sepsis, but should not delay care in order to obtain.**
- **Abnormally low temperatures increase mortality and are found often in geriatric patients.**
- Quantitative waveform capnography can be a reliable surrogate for lactate monitoring in detecting metabolic distress in sepsis patients. EtCO<sub>2</sub> < 25 mm Hg are associated with serum lactate levels > 4 mmol/L.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions** include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O<sub>2</sub> mask for the patient. This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected. A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions** include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions. This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions** include standard PPE plus airborne precautions plus contact precautions. This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. **SARS, SARS-CoV-2, COVID-19, MERS, Monkeypox**).
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen.
- Agency Medical Director may require contact of medical control prior to EMT / MR administering any medication.
- **Sepsis Screen:**
  - Agencies may use Adult / Pediatric Systemic Inflammatory Response Syndrome (SIRS) criteria or quickSOFA (qSOFA) criteria.
  - Receiving facility should be involved in determining Sepsis Screen utilized by EMS.



# Syncope

## History

- Cardiac history, stroke, seizure
- Occult blood loss (GI, ectopic)
- Females: LMP, vaginal bleeding
- Fluid loss: nausea, vomiting, diarrhea
- Past medical history
- Medications

## Signs and Symptoms

- Loss of consciousness with recovery
- Lightheadedness, dizziness
- Palpitations, slow or rapid pulse
- Pulse irregularity
- Decreased blood pressure

## Differential

- Vasovagal
- Orthostatic hypotension
- Cardiac syncope
- Micturition / Defecation syncope
- Psychiatric
- Stroke
- Hypoglycemia
- Seizure
- Shock (see Shock Protocol)
- Toxicological (Alcohol)
- Medication effect (hypertension)
- PE
- AAA

|   |   |
|---|---|
| Age Appropriate<br>Airway Protocol(s) AR 1, 2, 3, 5, 6<br><b>if indicated</b> |   |
|   | Blood Glucose Analysis Procedure  |
| <b>B</b>  | 12 Lead ECG Procedure   |
|   | IV or IO Access Protocol UP 6   |
| <b>P</b>  | Cardiac Monitor   |
|   | Altered Mental Status<br>Protocol UP 4<br><b>if indicated</b>   |
|   | Age Appropriate Cardiac<br>Protocol(s)<br><b>if indicated</b>   |
|   | Age Appropriate<br>Hypotension/ Shock<br>Protocol AM 5/ PM 3<br><b>if indicated</b>                           |
|   | Multiple Trauma Protocol TB 6<br>Spinal Motion Restriction<br>Procedure/ Protocol TB 8<br><b>if indicated</b> |

**Age Specific Blood Pressure  
indicating possible shock**

**Age 0 – 28 days: SBP < 60**  
**Ages ≥ 1 month: SBP < 70**  
**Age 1 – 9: SBP < 70 + (2x Age)**

**Ages 10 – 64: SBP < 90**  
**Ages ≥ 65: SBP < 110**

**All ages Shock Index:  
HR > SBP**

Serious Signs/ Symptoms  
Hypotension, poor  
perfusion, shock

NO

YES

|          |  |
|----------|--|
|          | IV or IO Access Protocol UP 6<br>Consider 2 Large Bore sites   |
| <b>A</b> | <b>Normal Saline / LR 500 mL Bolus</b><br>Repeat as needed<br>Titrate SPB ≥ 90 mmHg<br><b>Maximum 2 L</b>                        |
|          | <b>Peds: 20 mL/kg IV / IO</b><br>Repeat as needed<br>Titrate to Age Appropriate<br>SBP ≥ 70 + 2 x Age<br><b>Maximum 60 mL/kg</b> |

Exit to  
Age Appropriate  
Condition Appropriate  
Protocol(s)

**Notify Destination or  
Contact Medical Control**



# Syncope

## Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Syncope is both loss of consciousness and loss of postural/ muscle tone with collapse. Symptoms preceding the event are important in determining etiology.**
- **Syncope often is due to a benign process but can be an indication of serious underlying disease in both the adult and pediatric patient.**
- **Often patients with syncope are found normal on EMS evaluation. In general patients experiencing syncope require cardiac monitoring and emergency department evaluation.**
- **Differential should remain wide and include:**

|                           |                             |                    |                              |
|---------------------------|-----------------------------|--------------------|------------------------------|
| <b>Cardiac arrhythmia</b> | <b>Neurological problem</b> | <b>Choking</b>     | <b>Pulmonary embolism</b>    |
| <b>Hemorrhage</b>         | <b>Stroke</b>               | <b>Respiratory</b> | <b>Hypo or Hyperglycemia</b> |
| <b>GI Hemorrhage</b>      | <b>Seizure</b>              | <b>Sepsis</b>      |                              |
- **High-risk patients:**

|                                 |                                |
|---------------------------------|--------------------------------|
| <b>Age <math>\geq</math> 60</b> | <b>Syncope with exertion</b>   |
| <b>History of CHF</b>           | <b>Syncope with chest pain</b> |
| <b>Abnormal ECG</b>             | <b>Syncope with dyspnea</b>    |
- **Abdominal/ back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and/ or lower extremity pain or diminished pulses, especially in patients over 50 and/ or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 35, diabetics, and/ or women especially with upper abdominal complaints.**
- **Heart Rate: Tachycardia is one of the first clinical signs of dehydration, typically increases as dehydration becomes more severe.**
- **Syncope with no preceding symptoms or event may be associated with an arrhythmia.**
- **Assess for signs and symptoms of trauma if associated or questionable fall with syncope.**
- **Consider dysrhythmias, GI bleed, ectopic pregnancy, and seizure as possible causes of syncope.**
- **In general these patients should be transported: Patients who experience syncope associated with headache, neck pain, chest pain, abdominal pain, back pain, dyspnea, or dyspnea on exertion need prompt medical evaluation.**
- **More than 25% of geriatric syncope is cardiac dysrhythmia based.**



# Behavioral Health Crisis

## History


- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic alert tag
- Substance abuse / overdose
- Diabetes

## Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative violent
- Expression of suicidal / homicidal thoughts

## Differential

- Altered Mental Status
- Alcohol Intoxication
- Toxin / Substance abuse
- Medication effect / overdose / withdrawal
- Depression
- Bipolar (manic-depressive)
- Schizophrenia
- Anxiety disorders





# Behavioral Health Crisis

## Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Neurologic status**
- **Crew/ responders safety is the main priority. Call for assistance, stage, or withdraw from scene if necessary.**
- **Law Enforcement:**
  - **Any patient who is handcuffed or restrained by Law Enforcement and transported by EMS, must be accompanied by law enforcement during transport.**
  - **Patient should not be transported with upper extremities hand-cuffed behind back as this prevents proper assessment and could lead to injury.**
  - **Consider multidisciplinary coordination with law enforcement to approach verbal de-escalation, restraint, and/ or USP 6 Restraints: Therapeutic Take-down Procedure.**
- **Maintain high-index of suspicion for underlying medical or traumatic disorder causing or contributing to behavioral disturbance. Medical causes more likely in ages < 12 or > 40.**
- **General communications techniques**
  - **Ask Open-ended questions (questions that cannot be answered with a yes/no)**  
*“Tell me how we can help you?” “What caused you to call 911 today?”*
  - **Active listening (stay engaged, be able to summarize patient’s story, use your body language to convey listening)**  
*Eye contact, nodding your head, periodically repeating back part of patient’s story*
  - **Encouraging (remain positive, convey interest in patient’s crisis)**  
*“Tell me more about that...”*
  - **Clarifying questions (ask patient to rephrase or repeat if you don’t understand)**  
*“I’m not sure I understand, can you...?”*
  - **Emotional labeling (naming emotions patient is demonstrating, validating emotions)**  
*“You look upset.” “You seem angry.”*
  - **Conversational pause (okay to allow a period of silence for patient to process information)**
- **Behavioral health disturbance incidents are increasing and commonly involve the following:**

|  |  |
|--|--|
| <b>Substance misuse</b>                                | <b>Psychosis</b>                                   |
| <b>Depression/ Anxiety/ Stress Reactions / Bipolar</b> | <b>Schizophrenia or schizophrenia-like illness</b> |
- **Restraints:**
  - **All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
  - **Do not position or transport any restrained patient in such a way that could impact the patient’s respiratory or circulatory status (i.e. prone position)**
- **Maintain high-index of suspicion for medical, trauma, abuse, or neglect causes:**
  - **Hypoglycemia, hyperglycemia, overdose, substance abuse, hypoxia, head injury, shock, sepsis, stroke, etc.**
  - **Domestic violence, child or geriatric abuse/ neglect.**
- **Extrapyramidal reactions:**
  - **Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities. May present with contorted neck and trunk with difficult motor movements. Typically an adverse reaction to antipsychotic drugs like Haloperidol and may occur with your administration. When recognized, give **Diphenhydramine 50 mg IV / IO / IM / PO** in adults or **1 mg/kg IV / IO / IM / PO** in pediatrics, **Maximum 50 mg.****
- **May add page 3 to protocol for specific for local mental health and / or substance misuse resources or destinations.**



# Behavioral CIT Paramedic (Optional)

Complete EMS CIT Behavioral Health Data Sheet


Complete EMS CIT Behavioral Assessment Form

This is a continuation of the Evaluation and Screening for Mental Health and Substance Abuse Procedure

Alternative Destinations other than Hospital ED  
Call Access Line for all referrals:  
**Contact Number**

If appropriate, contact Mobile Crisis:  
**Contact Number**

- Blood/Vomit/Difficulty Breathing/Acute Medical Issue
- Blood Glucose <70 or >250 & symptomatic
- Head Trauma or fall in past 7 days
- In and out of consciousness
- Seizure activity in past 24 hours
- Unable to speak or walk
- On IVC or Emergency Custody
- Violent or Aggressive
- Provider can't see within 2 hours at alternative site
- Acute Withdrawal
- Hospital transport requested by patient
- BP ≥210/130 or ≥180/110 and symptomatic
- BAC > .35
- Oxygen Dependent




NO

- Suicidal (with plan and/or intent)
- Homicidal Ideation
- Psychosis (auditory/visual hallucinations or delusions)
- Request for Detox
- Med Request during 8:00 am and 3:00pm and client can not wait until next day for meds



NO

- No current suicidal ideation
- Anxiety/Panic
- Tearful crying with no suicidal plan or intent
- Med request in which EMS/Mobile Crisis able to triage and client is "ok" to go M-F for meds.




**Alternative Destinations / Crisis Providers For Centerpoint**

| County             |                    |                    |
|--------------------|--------------------|--------------------|
| Resource Agency    | Resource Agency    | Resource Agency    |
| Hours of Operation | Hours of Operation | Hours of Operation |




# Behavioral Agitation/ Sedation Guide





# Behavioral Hyperactive Delirium With Severe Agitation



**Call for help**  
**Call for additional resources**  
**Stage prior to arrival**  
 or  
**Withdraw from scene until safe**

|          |  |  |
|----------|--|--|
| <b>P</b> | <p><b>Ketamine 4 mg/kg IM</b><br/>         Maximum 400 mg <i>See Pearls</i></p> <p><b>Pediatrics:</b><br/>         2 mg/kg IM<br/>         Maximum 400 mg <i>See Pearls</i></p> <p>-----Or-----</p> <p><b>Midazolam 10 mg IM</b><br/>         +<br/> <b>Haldol 5mg IM</b><br/>         May repeat in 5-10 minutes as needed</p> <p>Or</p> <p><b>Geodon 20mg IM x1</b></p> <p><b>Pediatrics:</b><br/>         Midazolam 0.1 – 0.2 mg/kg IM<br/>         Max Single Dose 10mg<br/>         Repeat every 5 minutes as needed</p> <p><b>Haldol Max Total Dose 20 mg</b><br/> <b>Midazolam Max Total Dose 20 mg</b></p> |  |
|          | <p>IV or IO Access Protocol UP 6<br/>         Preferably 2 large bore</p>  |  |
| <b>A</b> | <p><b>Normal Saline / Lactated Ringers 1 L Bolus</b><br/>         Then 150 – 200 mL / hr<br/>         May repeat 500 mL Bolus<br/>         as needed<br/> <b>Maximum 2 L</b></p> <p><b>Peds: 20 – 60 mL/kg IV / IO</b><br/> <b>Maximum 60 mL/kg</b></p>  |  |
|          | <p>Restraint Physical Procedure USP 5<br/>         Monitor per restraint procedure<br/> <i>if indicated</i></p>  |  |
|          | <p>Restraint Therapeutic Take Down Procedure USP 6<br/> <i>if indicated</i></p>  |  |
|          | <p>Hyperthermia Protocol</p>   |  |
| <b>P</b> | <p>Cardiac Monitor</p>   |  |
|          | <p><b>Notify Destination or Contact Medical Control</b></p>  |  |

- General Guidance**
- Use of protocol is indicated when a medical emergency requires a combination of both physical restraint and chemical restraint in order to prevent imminent injury to a patient and/ or providers.
  - Use of this protocol requires medical judgement and consultation with Medical Control is not mandatory.
  - Non-medical personnel requests or opinions should not be used as a factor when implementing this protocol.



# Behavioral Hyperactive Delirium With Severe Agitation

All patient who have received medications per protocols UP18 & UP19 must be transported to the hospital or left in the care of a community paramedic.

Patients with a BARS Score of 7 who utilized this protocol must be transported to the hospital – No exceptions unless granted directly by Medical Director

We do not medicate for Jail. If IV/IO/IM medicine is given – the patient must be medically monitored and is not considered stable for jail. Releasing a sedated patient to law enforcement for jail will be considered abandonment. This population is at high risk for cardiac arrest post event.

Haldol + Midazolam have synergistic properties and it is recommended they are used together.

## Pearls

- **Ketamine for sedation purposes:**  
**Ketamine may be used in pediatric patients who fit within a Pediatric Medication/ Skill Resuscitation System product, ≤ 15 years of age, or ≤ 49 kg) with DIRECT ONLINE MEDICAL ORDER by the system MEDICAL DIRECTOR only.**
- **Hyperactive Delirium with Severe Agitation:**  
Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent/ bizarre behavior, insensitivity to pain, hyperthermia and increased strength.  
Potentially life-threatening and associated with use of physical control measures, including physical restraints.  
Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents.  
Alcohol or substance withdrawal as well as head trauma may also contribute to the condition.
- **Restraint use:**  
Physical restraints are not contraindicated in agitated or excited delirium, but you must use caution. Once sedated, prevent patient from continued struggle, which can worsen metabolic condition. Prevent patient from assuming a prone position for prolonged period, move to supine position as quickly as possible.  
Team approach for sedation and Restraint Therapeutic Take Down Procedure USP-6:
  - 1 provider for each limb.
  - 1 provider to lead restraint, maintain airway and control head.
  - 1 Provider to administer medication.Do not position prone or prone with restraints, as this can impede respiration and ventilation.
- Hyperthermia: Assess for and treat hyperthermia.



# Sickle Cell Crisis

## History

- Past medical history
- Medications
- Recent illness
- Prior pain crisis location
- Pain regimen at home

## Signs and Symptoms

- Pain
- One sided paralysis / weakness
- Difficulty walking / speaking
- Sudden vision changes
- Unexplained numbness
- Severe headache
- Fever
- SOB
- Chest Pain
- Abdominal Pain
- Pallor

## Differential

- Sickle Celle Pain Crisis
- Aplastic Crisis
- Acute Chest Syndrome
- Alcohol / drug use
- Toxic ingestion
- Seizure
- Stroke
- Altered baseline mental status
- Sepsis
- Pneumonia

Apply Hot Packs to affected areas: especially joints and areas of increased pain

Provide emotional support  
Calm and continual reassurance

|          |   |
|----------|---|
| <b>B</b> | Blood Glucose Analysis as needed<br>Assess Pain Severity<br>12 Lead ECG Procedure |
|          | IV or IO Access Protocol UP 6   |
| <b>P</b> | Cardiac Monitor   |
|          | Altered Mental Status<br>Protocol UP 4<br><i>if indicated</i>                     |
|          | Manage Airway as appropriate<br>Protocol AR 1 & AR 5<br><i>if indicated</i>       |
|          | Suspected Stroke<br>Protocol UP 14<br><i>if indicated</i>                         |
|          | Suspected Seizure<br>Protocol UP 13<br><i>if indicated</i>                        |
|          | Hypotension/ Shock<br>Protocol AM 5 & PM 3<br><i>if indicated</i>                 |

Fluids indicated if signs of shock or hypotensive

**Sickle Cell Crisis**  
- Acute Chest Syndrome  
- Abdominal Crisis  
- Joint Crisis

YES

Pain Control  
Protocol UP 11

**Notify Destination or Contact Medical Control**



# Sickle Cell Crisis

## PEARLS

Patterns of an acute sickle cell crisis are now recognizable. They are based on the part of the body where the crisis occurs.

Is this their typical pain crisis? If not, what is different about it?

Any fever, SOB, pleuritic chest pain?

### **Acute chest syndrome:**

Sudden acute chest pain with coughing up of blood can occur. Low-grade fevers can be present. The person is usually short of breath. If a cough is present, it often

is nonproductive. Acute chest syndrome is common in a young person with sickle cell disease.

Chronic (long-term) sickle cell lung disease develops over time because the acute and subacute lung crisis leads to scarred lungs as well as other problems.

### **Abdominal crisis:**

The pain associated with the abdominal crisis of sickle cell disease is constant and sudden. It becomes unrelenting. The pain may or may not be localized to any one area of the abdomen. Nausea, vomiting, and diarrhea may or may not occur.

### **Joint crisis:**

Acute and painful joint crisis may develop without a significant traumatic history. Its focus is either in a single joint or in multiple joints. Often the connecting bony parts of the joint are painful. Range of motion is often restricted because of the pain. Avascular necrosis of the hips can occur, causing permanent damage.

### **Stroke:**

Many sickle cell patients can present with strokes at a younger age than average. Ensure hospital pre-notification indicating it is a sickle cell patient as the standard of care for a Sickle Cell Stroke can involve exchange therapy as opposed to other treatments such as tPA.