



# Adult Polymorphic Tachycardia WIDE ( $\geq 0.12$ sec) Torsades de pointes

### History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

### Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

### Differential

- Cardiac arrest
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose

**Assess tachycardia in context of clinical condition  
Identify and treat underlying cause of tachycardia**

**Unstable / Serious Signs and Symptoms  
HR Typically  $\geq 150$**   
Hypotension, Acute AMS, Ischemic Chest Pain,  
Acute CHF, Seizures, Syncope, or Shock  
secondary to tachycardia

|          |   |
|----------|---|
| <b>P</b> | Defibrillation Procedure  |
|          | <b>Consider Sedation Prior to Defibrillation</b><br><b>Midazolam 2 – 5 mg IV / IO / IN</b><br>May repeat as needed<br><b>Maximum 10 mg</b><br>Consider Ketamine 1mg/kg IV / IM<br><b>Wide and Irregular: 200 – 360J</b> |
|          | <b>Polymorphic QRS (Not-Synchronized)</b><br><br><i>May repeat and increase dose with subsequent cardioversion attempts</i>   |

|          |                               |
|----------|-------------------------------|
| <b>B</b> | 12 Lead ECG Procedure         |
| <b>P</b> | Cardiac Monitor               |
|          | IV or IO Access Protocol UP 6 |

**Pulse Present?**

NO  
Exit to  
Cardiac Arrest  
Protocol AC 3

YES → **PROCEED**

QT Interval < 500 msec

|          |  |
|----------|--|
| <b>P</b> | <b>Amiodarone 150 mg in<br/>100 mL of D5W IV / IO</b><br>Infuse over 10 minutes<br>May repeat if tachycardia recurs or persists                              |
|          | Or<br><br><b>Lidocaine<br/>1 – 1.5 mg/kg IV / IO</b><br><br>May repeat if refractory<br><b>Lidocaine<br/>1.5 mg/kg IV / IO</b><br><br><b>Maximum 3 mg/kg</b> |
|          | Monitor and Reassess   |

QT Interval > 500 msec

|          |   |
|----------|---|
| <b>P</b> | <i>Consider</i><br><b>Magnesium 2 g IV / IO</b><br><br>May repeat<br><br><b>Maximum 4 g</b> |
|          | Monitor and Reassess  |

**Polymorphic QRS:**  
• QRS complexes in a single lead will change shape from complex to complex.

**Notify Destination or Contact Medical Control**



# Adult Polymorphic Tachycardia WIDE ( $\geq 0.12$ sec) Torsades de pointes

Do not use amiodarone for individuals with polymorphic VT associated with a prolonged QT interval because this may worsen the patient's condition. Administer Magnesium, loading dose 2 Gm IV/IO for polymorphic VT.

## Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
  - **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
  - **12-Lead ECG:**
    - 12 Lead ECG not necessary to diagnose and treat
    - Obtain when patient is stable and/or following rhythm conversion.
  - **Monomorphic QRS:**
    - All QRS complexes in a single lead are similar in shape.
  - **Polymorphic QRS:**
    - QRS complexes in a single lead will change shape from complex to complex.
  - **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
  - **Unstable condition**
    - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
    - If at any point patient becomes unstable move to unstable arm in algorithm.
  - **Symptomatic condition**
    - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
    - Symptomatic tachycardia usually occurs at rates  $\geq 150$  beats per minute. Patients symptomatic with heart rates  $< 150$  likely have impaired cardiac function such as CHF.
  - **Serious Signs / Symptoms:**
    - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
  - **Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.**
  - **Typical sinus tachycardia is in the range of 100 to (220 – patients age) beats per minute.**
  - **If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW), DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.**
  - **Polymorphic / Irregular Tachycardia:**
    - This situation is usually unstable and immediate defibrillation is warranted.
    - If QT length is known, use for decision-making. Prolonged QT length defined as  $> 500$  msec.
    - QT length  $< 500$  msec:
      - Arrhythmia more likely related to ischemia or infarction and Magnesium not likely helpful.
      - May quickly deteriorate into Ventricular Fibrillation.
      - Even when terminated by defibrillation, may recur, so follow with medication therapy.
    - QT prolongation  $> 500$  msec:
      - Magnesium more likely to be helpful.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.