



# Pediatric Tachycardia

## Wide Complex (> 0.09 sec)

### History


- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

### Signs and Symptoms

- Heart Rate: Child > 180/bpm  
Infant > 220/bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

### Differential

- Heart disease (Congenital)
- Hypothermia/ Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety/ Pain/ Emotional stress
- Fever/ Infection/ Sepsis
- Hypoxia, Hypoglycemia
- Medication/ Toxin/ Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax





# Pediatric Tachycardia

## Wide Complex (> 0.09 sec)

If Fever 101F or above - likely Sepsis -- treat underlying problem not tachycardia !

If Heartrate increases with Adenosine then likely WPW – No further Adenosine. Absolutely No Cardizem – Contact Med Control

### Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Neuro**
- **Monomorphic QRS:**
  - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
  - QRS complexes in a single lead will change from complex to complex.
- Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.
- Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.
- **12-Lead ECG:**
  - 12-Lead ECG is not necessary to diagnose and treat arrhythmia. A single lead ECG is often all that is needed.
  - Obtain 12-Lead when patient is stable and/ or following a rhythm conversion.
  - When administering adenosine, obtaining a continuous 12-Lead can be helpful later to physicians.
- **Unstable condition:**
  - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
  - If at any point patient becomes unstable move to unstable arm in algorithm
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.
- **Serious Signs and Symptoms:**
  - Respiratory distress/ failure.
  - Signs of shock/ poor perfusion with or without hypotension.
  - AMS
  - Sudden collapse with rapid, weak pulse
- **Serious Signs and Symptoms:**
  - Respiratory distress/ failure.
  - Signs of shock/ poor perfusion with or without hypotension.
  - AMS
  - Sudden collapse with rapid, weak pulse
- **Wide Complex Tachycardia (≥ 0.09 seconds):**
  - SVT with aberrancy.
  - VT: Uncommon in children. Rates may vary from near normal to > 200/ minute.
  - Most children with VT have underlying heart disease / cardiac surgery/ long QT syndrome/ cardiomyopathy.
  - **Amiodarone 5 mg / kg over 20 – 60 minutes or Procainamide 15 mg / kg over 30 – 60 minutes IV / IO** are recommended agents. They should not be administered together. Consultation with Medical Control is advised when these agents are considered.
- **Torsade's de Pointes/ Polymorphic (multiple shaped) Tachycardia:**
  - Rate is typically 150 to 250 beats/ minute.
  - Associated with long QT syndrome, hypomagnesaemia, hypokalemia, many cardiac drugs.
  - May quickly deteriorate to VT.
  - Separating the child from the caregiver may worsen the child's clinical condition.
- Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.
- Continuous pulse oximetry is required for all SVT patients if available.